

A service of the National Library of Medicine
and the National Institutes of Health

www.pubmed.gov

My NCBI

[\[Sign In\]](#) [\[Regis\]](#)[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)[Scol](#)Search **PubMed** for ☒ Limits ☐ Preview/Index ☐ History ☐ Clipboard ☐ DetailsField: **Pagination, Limits: Publication Date from 1995 to 1995**Display **Abstract** Show **20** Sort by Send to All: 1 Review: 0 [About Entrez](#)
[NCBI Toolbar](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorials](#)[New/Noteworthy](#)[E-Utilities](#)[PubMed Services](#)[Journals Database](#)[MeSH Database](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[Special Queries](#)[LinkOut](#)[My NCBI](#)[Related Resources](#)[Order Documents](#)[NLM Mobile](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)☐ **1: Microbiology.** 1995 Feb;141 (Pt 2):281-90.[Related Articles, Links](#)**Genes encoding xylan and beta-glucan hydrolysing enzymes in *Bacillus subtilis*: characterization, mapping and construction of strains deficient in lichenase, cellulase and xylanase.****Wolf M, Geczi A, Simon O, Borriß R.**

Institut für Biologie, Humboldt Universität, Berlin, Germany.

The gene encoding extracellular xylanase (xynA) was amplified as a 770 bp DNA fragment from *Bacillus subtilis* 168 chromosomal DNA by PCR. The genes encoding endo-beta-1,4-glucanase (eglS) and endo-beta-1,3-1,4-glucanase (bglS) were isolated from a genomic library of *B. subtilis* 168. The sequences of xynA and eglS were identical to those of the xylanase and cellulase genes from *B. subtilis* PAP115. Integrative plasmids containing DNA fragments with deletions in the coding region of the genes were constructed and used to replace the chromosomal eglS, bglS and xynA genes of *B. subtilis* 168. Strains without any detectable activity against xylan (Xyn-), carboxymethylcellulose (Egl-) or mixed linked beta-1,3-1,4-glucan (Egl- Bgl-) were obtained. The genes were mapped at 170 degrees (eglS), 175 degrees (xynA) and 340 degrees (bglS) on the *B. subtilis* chromosome.

PMID: 7704256 [PubMed - indexed for MEDLINE]

Display **Abstract** Show **20** Sort by Send to [Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Feb 24 2006 04:49:50